ELSEVIER

www.elsevier.nl/locate/jorganchem

Journal of Organometallic Chemistry 598 (2000) 419–420

Journal of Organometallic Chemistry

Author Index of Volume 598

Abri, A., 222
Adams, H., 1
Ader, B., 24
Akhmedov, N.G., 387
Antiñolo, A., 167
Arques, A., 329
Asadi, M.G., 222
Aucar, G.A., 193
Auñón, D., 329
Avtomonov, E.V., 403

Bardin, V.V., 127
Bautista, D., 329
Bedford, R.B., 20
Bentabet, E., 365
Bernardini, R., 174
Bickley, J.F., 409
Blacque, O., 365
Botek, E.L., 193
Boyer, P., 365
Brauer, D.J., 116
Broussier, R., 365
Bruhn, C., 286
Burlakov, V.V., 243
Butler, I.R., 103
Byrne, L.T., 28

Carvalho, M.F.N.N., 318
Castro-Juiz, S., 71
Caubet, A., 87
Cazin, C.S.J., 20
Cea-Olivares, R., 160
Čermák, J., 318
Chan, K.S., 80
Chao, Y.-C., 13
Charrue, P., 187
Chen, B.-H., 348
Chen, W., 409
Chen, Z.-N., 63
Cheung, K.-K., 55
Clark, A.M., 262
Clegg, W., 248
Coles, S.J., 103
Cory, M.G., 193
Cruz-Almanza, R., 160

Dakternieks, D., 49
Day, C.S., 150
del Hierro, I., 167
de Montauzon, D., 215
Dong, Y.-B., 55

Elsegood, M.R.J., 248
Englich, U., 202
Ephritikhine, M., 313

Fajardo, M., 167
Fang, M., 42
Fargeas, V., 187
Favresse, F., 187
Fernández, A., 1, 71
Fernández, J.J., 1, 71
Ferrer, M., 215
Fiedler, J., 136
Font-Bardía, M., 87, 215
Foos, E.E., 182
Franke, H., 127
Fritzen, P., 127
Frohn, H.-J., 127
Fu, E., 42
Furin, G.G., 243

Galvão, A.M., 318
Gao, C., 42
García, M.M., 254
García y García, P., 160
García-Yuste, S., 167
Gautheron, B., 365
Gerisch, M., 286
Görls, H., 108
Graschy, S., 202
Gross, T., 395

Haukka, M., 235
Häusler, T., 116
Heikkinen, V., 235
Hengge, E., 202
Herberhold, M., 142
Hermann, U., 202
Hihara, G., 276
Hoffmann, D., 395
Hoffmann, T., 286
Horrocks, B.R., 248
Hos, J.P., 28
Houlton, A., 248
Huang, J., 339
Huang, S.-P.D., 374
Hurley, A.L., 150
Hursthouse, M.B., 103
Hynes, R.C., 276

Isaac, C.J., 248
Israel, G., 286

Jia, G., 228
Jouet, R.J., 182

Kaim, W., 136
Kang, B.-S., 63
Karlov, S.S., 387
Kastner, G., 381
Kempe, R., 395
Kiae, H., 222
Klimova, E.I., 254
Klimova, T., 254
Knoesen, O., 108
Ko, B.-T., 13
Kooijman, H., 24
Koskinen, A.M.P., 235
Koutsantonis, G.A., 28
Kubicki, M.M., 365

Lahoz, F.J., 329
Laitinen, R.H., 235
Lance, M., 313
Lazzaroni, R., 174
Le Borgne, T., 313
Lebret, B., 187
Lebuis, A.-M., 276
Lee, G.-H., 353
Leggett, G., 36
Leng, M., 339
Letov, A.V., 243
Liao, M.-S., 374
Licence, P., 103
Lin, C.-C., 13
Liu, C.-M., 348
Liu, W.-Y., 348
Li, Z., 339
López, C., 87
López-Solera, I., 167
López-Torres, M., 1, 71
Lorberth, J., 387
Lotz, S., 108

Mak, T.C.W., 80
Marschner, C., 202
Martínez, F., 329
Mayr, A., 55
Ma, Y.-X., 348
Mechtler, C., 202
Mellet, P., 365
Milius, W., 142
Minacheva, M.K., 243
Mohammadpour, A., 222
Molina, P., 329
Motevalli, M., 36
Müller, F.W., 286

Nierlich, M., 313

O'Connell, J., 49
Oehme, H., 395
Okuda, J., 179
Onyszchuk, M., 276
Ortigueira, J.M., 1, 71
Otero, A., 167

Passler, T., 292, 304
Peng, S.-M., 353
Pereira, T., 1

PII: S0022-328X(00)00052-8

Petz, W., 403
Pinter, E., 202
Piteau, M., 187
Podlaha, J., 318
Polborn, K., 292
Pombeiro, A.J.L., 318
Ponomarev, S.V., 211
Price, C., 248
Pursiainen, J., 235

Qian, Y., 339
Quintard, J.-P., 187

Ramirez, L.R., 254
Reuter, H., 381
Rickard, C.E.F., 262
Rivière-Baudet, M., 276
Rodríguez, A., 167
Roßenbach, S., 116
Roper, W.R., 262
Rosenthal, U., 243
Rossell, O., 215
Ruhlandt-Senge, K., 202

Šabata, S., 318
Safa, K.D., 222
Salvadori, P., 174
Scheiring, T., 136
Schenk, S., 116
Seco, M., 215
Seip, M.A., 387
Sheldrick, W.S., 116
Shen, Q., 359
Shur, V.B., 243
Shutov, P.L., 387
Sinnema, P.-J., 179
Skelton, B.W., 28
Solans, X., 87, 215
Soler, M., 215
Spaniol, T.P., 179
Spee, M.P.R., 24
Spek, A.L., 24
Steenwinkel, P., 24
Steinborn, D., 286
Steiner, A., 409
Stelzer, O., 116
Strunkina, L.I., 243
Suárez, A., 1, 71
Sullivan, A.C., 36

Teijido, B., 71
Tepper, M., 116
Tiekink, E.R.T., 49
Toledano, C.A., 254
Tong, Y.-X., 63
Toscano, R.-A., 160
Toscano, R.A., 254
Troyanov, S.I., 243
Tse, M.K., 80

Uccello-Barretta, G., 174
Uhlig, F., 202

van Koten, G., 24
Velasco, M.D., 329
Vila, J.M., 1, 71
Vilaplana, M.J., 329
Vojtíšek, P., 318

Wagner, S., 292, 304
Wang, G., 42
Wang, Y., 359
Welker, M.E., 150
Weller, F., 403
Wells, R.L., 182
Wharf, I., 276
White, A.H., 28
White, P.S., 182
Wiberg, N., 292, 304
Wong, W.T., 228
Wrackmeyer, B., 211
Wright, L.J., 262
Wu, C., 42
Wu, D.-D., 80
Wu, X.-L., 348

Xiao, J., 409
Xue, F., 359
Xue, P., 42
Xu, L., 409

Yang, C.-C., 353
Yang, L., 55
Yan, H., 142
Yao, J., 228
Yao, T., 339
Yeh, W.-Y., 353
Yu, K., 359
Yu, K.-B., 63

Zaitseva, G.S., 387
Zerner, M.C., 193
Zhang, H.-X., 63
Zhou, X., 80

ELSEVIER

www.elsevier.nl/locate/jorganchem

Journal of Organometallic Chemistry 598 (2000) 421–429

Journal of Organometallic Chemistry

Subject Index of Volume 598

Acetylene complexes
Interaction of bis(trimethylsilyl)acetylene complex of titanocene with tris(pentafluorophenyl)borane. Synthesis and structure of a new type of zwitterionic metallocene (η^5-C_5H_5){η^5-[$C_5H_4B(C_6F_5)_3$]}Ti, 243

Acetyl ligand
Geometrical and electronic structures of the acetyl complex Re(bpy)(CO)$_3$($COCH_3$) and of [M(bpy)(CO)$_4$](OTf), M = Mn,Re, 136

Aggregation
Syntheses of cationic four-coordinated aluminum complexes: counter-ion induced aggregation through the N–H···I hydrogen bonds, 13

Alkenes
Synthesis and decomposition of bis(alkoxyethynyl)dimethyltin, 211

Alkenyl
Alkyne insertion reactions of [RuH(κ^2-S_2CNEt_2)(CO)(PPh_3)$_2$]: synthesis of alkenyl, alkynyl and enynyl complexes, 20

Alkyl and acyl oxo niobocene
Synthesis and reactivity of new oxo alkyl or oxo acyl niobocene complexes and crystal structure of $Cp'_2Nb(=O)(OC(O)CF_3)$ ($Cp' = \eta^5$-$C_5H_4SiMe_3$), 167

Alkylation
Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254

Alkynes
Alkyne insertion reactions of [RuH(κ^2-S_2CNEt_2)(CO)(PPh_3)$_2$]: synthesis of alkenyl, alkynyl and enynyl complexes, 20
Synthesis and decomposition of bis(alkoxyethynyl)dimethyltin, 211

Alkynyl complexes
Reactions of transition-metal η^1-propargyl complexes with sulfur dioxide and sulfur trioxide: transition-metal–carbon bond cleaving reactions of the cycloadducts which yield 1,2-oxathiole-2-oxides and -2,2-dioxides, 150

Aluminium
Isolation of new polynuclear aluminium amides; trinuclear [Al-Cl(MeNCH$_2$CH$_2$NMe)$_2$(Al(NMe$_2$)Cl)(Al(NMe$_2$)$_2$)] and fluxional adamantane-like [$Al_4Cl_2(NMe_2)_6$(NMe)O], 36

Aluminum
Syntheses of cationic four-coordinated aluminum complexes: counter-ion induced aggregation through the N–H···I hydrogen bonds, 13

Amides
Isolation of new polynuclear aluminium amides; trinuclear [Al-Cl(MeNCH$_2$CH$_2$NMe)$_2$(Al(NMe$_2$)Cl)(Al(NMe$_2$)$_2$)] and fluxional adamantane-like [$Al_4Cl_2(NMe_2)_6$(NMe)O], 36

Amido ligand
Non-bridged amido cyclopentadienyl complexes of titanium: synthesis, characterization, and olefin polymerization catalysis, 179

β-Arylvinyl azides
P,N-Heterodifunctional ligands by selective Staüdinger reaction of β-substituted vinylazides with (*Z*)-1,2-bis(diphenylphosphanyl)ethene and formation of cyclometalled complexes of palladium(II) of these ligands — crystal and molecular structure of a new chiral cyclometallaphosphoraniminophosphane of palladium(II), 329

Atomic parameters
Implementation of the IPPP–CLOPPA–INDO/S method for the study of indirect nuclear spin coupling constants and its application to molecules containing tin nuclei, 193

Basicity
2-Thioanisyldichlorophosphine, new starting material for the preparation of multidentate phosphine ligands: syntheses and characterization of derivatives of 2-anisyl- and 2-thioanisyldichlorophosphines, 235

Benzyl cation
Non-bridged amido cyclopentadienyl complexes of titanium: synthesis, characterization, and olefin polymerization catalysis, 179

2,2′-Bipyridine
Development of a T-joint for covalent molecular construction based on 2,2′-bipyridine and phenanthroline isocyanide metal complexes, 55

Bis(diphenylphosphino)ferrocene
Ferrocenyl ligands in mixed valence high-oxidation state rhenium complexes, 108

Bis(ferrocenylmethyl)ethylenediamine
Ferrocenyl ligands in mixed valence high-oxidation state rhenium complexes, 108

Boranes
(Fluoroorgano)fluoroboranes and -fluoroborates I: synthesis and spectroscopic characterization of potassium fluoroaryltrifluoroborates and fluoroaryldifluoroboranes, 127

Borates
(Fluoroorgano)fluoroboranes and -fluoroborates I: synthesis and spectroscopic characterization of potassium fluoroaryltrifluoroborates and fluoroaryldifluoroboranes, 127

Boronic acid
(Fluoroorgano)fluoroboranes and -fluoroborates I: synthesis and spectroscopic characterization of potassium fluoroaryltrifluoroborates and fluoroaryldifluoroboranes, 127

Bulky ligand
Mesitylpseudohalogermanes $Mes_2Ge(CN)_2$ and Mes_3GeX (X = CN, NCS, N_3, NCO and OH; Mes = 2,4,6-trimethylphenyl): syntheses, crystal and molecular structures, 276

Cage ion
Monoorganotin–polyoxometal compounds: Part 1. Synthesis and crystal structure of [(^{t}BuSn)$_{11}$(V^{IV}O)$_{12}$(OH)$_6$]Cl·2DMF·H_2O. The first example in a series of monoorganotin–polyoxometal compounds with icosahedral metal arrangement and various contents of tin and vanadium, 381

PII: S0022-328X(00)00053-X

Carbene complexes
Carbonyl insertion into zirconium–nitrogen bonds; synthesis and X-ray structure of a carbene complex composed of $[Zr(NMe_2)_4]_2$ and three $Fe(CO)_5$ units containing Fe→Zr donor–acceptor interactions, 403
Carbenes
Fluoroalkylated *N*-heterocyclic carbene complexes of palladium, 409
Carbenoid intermediate
Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254
Carbide
Reactions of metalloalkynes: Part 6. Synthesis of open tetrametallic clusters. Reaction of $[Ru(CO)_4(ethene)]$ with ethyne-1,2-diyl compounds. Single crystal X-ray structural determinations of $[Ru_4(\mu_4\text{-}CC)(\eta\text{-}C_5H_4R)_2(\mu\text{-}CO)_2(CO)_8]$ (R = H, Me), 28
Carbocation
Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254
Carbonyl complexes
Geometrical and electronic structures of the acetyl complex $Re(bpy)(CO)_3(COCH_3)$ and of $[M(bpy)(CO)_4](OTf)$, M = Mn,Re, 136
Catalysis
Synthesis of the diphenylacetylene-based, tetra-amine ligand 1,2-bis(3,5-bis[(dimethylamino)methyl]phenyl)acetylene by palladium-catalysed cross-coupling: isolation and crystal structure of the catalyst *trans*-(3,5-bis[(dimethylamino)-methyl]phenyl)bis(triphenylphosphine)palladium(II) iodide, 24
Cation
Syntheses of cationic four-coordinated aluminum complexes: counter-ion induced aggregation through the N–H···I hydrogen bonds, 13
C–H activation
Diverse reactivity of rhodium β-(tetraphenyl)tetraphenyl porphyrin chlorides with benzonitrile: formation of Rh porphyrin arene and imine complexes, 80
Chiral derivatizing agent
Use of *cis*-dichloro[(*S*)-α-methylbenzylamine](ethylene)Pt(II) as a chiral derivatizing agent for the determination by ^{195}Pt-NMR of the enantiomeric composition of unsaturated ethers or alcohols having a quaternary chiral carbon atom, 174
Chromium group
Syntheses, characterization and structures of chromium group carbonyl complexes containing a multifunctional $Ph_2P(o\text{-}C_6H_4)CH{=}N(CH_2)_2(o\text{-}C_6H_4N)$ ligand, 353
Closed-shell interactions
Crystal structures and spectroscopic properties of platina-β-diketones — a new mode of stacking in platinum complexes, 286
Cluster
Reactions of metalloalkynes: Part 6. Synthesis of open tetrametallic clusters. Reaction of $[Ru(CO)_4(ethene)]$ with ethyne-1,2-diyl compounds. Single crystal X-ray structural determinations of $[Ru_4(\mu_4\text{-}CC)(\eta\text{-}C_5H_4R)_2(\mu\text{-}CO)_2(CO)_8]$ (R = H, Me), 28
Clusters
Synthesis, characterization and solution studies on heteronuclear iron–copper clusters containing diphosphine ligands: crystal structure of the dicluster $[\{Fe_2(\mu\text{-}CO)(CO)_6(\mu\text{-}PPh_2)\text{-}Cu\}_2(dppp)]$, 215
Coordination
Synthesis and properties of some novel ferrocene macrocyclic dioxopolyamines, 42
Copper
Synthesis, characterization and solution studies on heteronuclear iron–copper clusters containing diphosphine ligands: crystal structure of the dicluster $[\{Fe_2(\mu\text{-}CO)(CO)_6(\mu\text{-}PPh_2)\text{-}Cu\}_2(dppp)]$, 215
Crystal field effects
A theoretical study of gas-phase and solid-state $Hg(CF_3)_2$, 374
Crystal structure
Carbonyl insertion into zirconium–nitrogen bonds; synthesis and X-ray structure of a carbene complex composed of $[Zr(NMe_2)_4]_2$ and three $Fe(CO)_5$ units containing Fe→Zr donor–acceptor interactions, 403
Crystal structures and spectroscopic properties of platina-β-diketones — a new mode of stacking in platinum complexes, 286
Geometrical and electronic structures of the acetyl complex $Re(bpy)(CO)_3(COCH_3)$ and of $[M(bpy)(CO)_4](OTf)$, M = Mn,Re, 136
Influence of phenyl ring substituents in the cyclometallation of Schiff base ligands: crystal and molecular structures of $[Pd\text{-}\{3,4\text{-}(OCH_2O)C_6H_2C(H){=}N(Cy)\text{-}C2,N\}(\mu\text{-}O_2CMe)]_2$ and $[Pd\text{-}\{3,4\text{-}(OCH_2CH_2O)C_6H_2C(H){=}N(Cy)\text{-}C6,N\}(\mu\text{-}O_2CMe)]_2$, 71
Monoorganotin–polyoxometal compounds: Part 1. Synthesis and crystal structure of $[(^tPrSn)_{11}(V^{IV}O)O_{14}(OH)_6]Cl\cdot 2DMF\cdot H_2O$. The first example in a series of monoorganotin–polyoxometal compounds with icosahedral metal arrangement and various contents of tin and vanadium, 381
Syntheses of titanium μ-arylimido and μ-pyridylimido complexes bearing (un)substituted cyclopentadienyl ligand, 339
Synthesis and crystal structure of $[U(\eta\text{-}C_8H_8)]_2[\mu\text{-}\eta^4\text{:}\eta^4\text{-}HN(CH_2)_3N(CH_2)_2N(CH_2)_3NH]$, a dinuclear compound with a bridging tetra-amide ligand, 313
Crystal structures
Ferrocenyl ligands in mixed valence high-oxidation state rhenium complexes, 108
Mesitylpseudohalogermanes $Mes_2Ge(CN)_2$ and Mes_3GeX (X = CN, NCS, N_3, NCO and OH; Mes = 2,4,6-trimethylphenyl): syntheses, crystal and molecular structures, 276
Cyano-bridged complex
Cyano-bridged extended heteronuclear supramolecular architectures with hexacyanoferrates(II) as building blocks, 63
Cyclic voltammetry
Synthesis and properties of some novel ferrocene macrocyclic dioxopolyamines, 42
[3 + 2] Cycloaddition
Reactions of transition-metal η^1-propargyl complexes with sulfur dioxide and sulfur trioxide: transition-metal–carbon bond cleaving reactions of the cycloadducts which yield 1,2-oxathiole-2-oxides and -2,2-dioxides, 150
[3 + 2]-Cycloadditionen
Addukte der Ethene $Me_2E{-}C(SiMe_3)_2$ (E = Si, Ge, Sn) mit LiR und RN_3: Wie rasch bilden sie sich?, 304
Dimere der Ethene $Me_2E{-}C(SiMe_3)_2$ (E = Si, Ge, Sn): Auf welchem Wege entstehen sie aus $Me_2EX{-}CM(SiMe_3)_2$? Wie sind sie strukturiert?, 292
Cyclometallated
Reactivity of cyclometallated semicarbazone complexes of Pd(II): crystal and molecular structures of $[Pd\{C_6H_4C(Et){=}NN(H)C(=O)NH_2\}(PPh_3)][ClO_4]$ and $[Pd\{C_6H_4C(Et){=}NN(H)C(=O)NH_2\}\{(Ph_2PCH_2CH_2)_2PPh\text{-}P,P,P\}][Cl]$, 1
Cyclometallation
Influence of phenyl ring substituents in the cyclometallation of Schiff base ligands: crystal and molecular structures of $[Pd\text{-}\{3,4\text{-}(OCH_2O)C_6H_2C(H){=}N(Cy)\text{-}C2,N\}(\mu\text{-}O_2CMe)]_2$

and [Pd-{3,4-(OCH$_2$CH$_2$O)C$_6$H$_2$C(H)=N(Cy)-*C*6,*N*}(μ-O$_2$-CMe)]$_2$, 71
Cyclometalled complexes
P,*N*-Heterodifunctional ligands by selective Staüdinger reaction of β-substituted vinylazides with (*Z*)-1,2-bis(diphenylphosphanyl)ethene and formation of cyclometalled complexes of palladium(II) of these ligands — crystal and molecular structure of a new chiral cyclometallaphosphoraniminophosphane of palladium(II), 329
Cyclopalladation
Platinum(II) and palladium(II) compounds derived from [(η^5-C$_5$H$_5$)Fe{(η^5-C$_5$H$_4$)–CH=N–CH$_2$–CH$_2$–OH}], 87
Cyclopropane
Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254
Cyclopropene
Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254
Cyclosilanes
On some reactions of undecamethylcyclohexasilanyl-substituted silanes, 202

Density-functional calculations
A theoretical study of gas-phase and solid-state Hg(CF$_3$)$_2$, 374
Deprotonation
Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254
Dichlorophosphines
2-Thioanisyldichlorophosphine, new starting material for the preparation of multidentate phosphine ligands: syntheses and characterization of derivatives of 2-anisyl- and 2-thioanisyldichlorophosphines, 235
Diene adduct
Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254
Diethynylbenzene
Alkyne insertion reactions of [RuH(κ^2-S$_2$CNEt$_2$)(CO)(PPh$_3$)$_2$]: synthesis of alkenyl, alkynyl and enynyl complexes, 20
Dinuclear
Synthesis and crystal structure of [U(η-C$_8$H$_8$)$_2$]$_2$[μ-η^4:η^4-HN(CH$_2$)$_3$N(CH$_2$)$_2$N(CH$_2$)$_3$NH], a dinuclear compound with a bridging tetra-amide ligand, 313
Diphenylamide complex
Heteroleptic diphenylamido complexes of ytterbium: syntheses and molecular structures of (MeC$_5$H$_4$)$_2$YbNPh$_2$(THF) and (C$_5$Me$_5$)$_2$YbNPh$_2$, 359
Diphosphine
Synthesis, characterization and solution studies on heteronuclear iron–copper clusters containing diphosphine ligands: crystal structure of the dicluster [{Fe$_2$(μ-CO)(CO)$_6$(μ-PPh$_2$)Cu}$_2$(dppp)], 215
1,2-Disilacyclobutanes
Head-to-head versus head-to-tail dimerizations of transient silenes — the generation and dimerization behavior of 2-(2-dimethylaminoaryl)-1,1-bis(trimethylsilyl)silenes, 395
1,3-Disilacyclobutanes
Head-to-head versus head-to-tail dimerizations of transient silenes — the generation and dimerization behavior of 2-(2-dimethylaminoaryl)-1,1-bis(trimethylsilyl)silenes, 395
Dithiophosphates
Synthesis, characterization and X-ray structure of stannocanes substituted with a cyclic dithiophosphate ligand X(CH$_2$-CH$_2$S)$_2$Sn^{n}Bu [S$_2$P(OCH$_2$C(Et$_2$)CH$_2$O)], (X = O, S): a study about the conformational tendencies and the relationship with the anomeric effect of the stannocane rings, 160
Diverse reactivity
Diverse reactivity of rhodium β-(tetraphenyl)tetraphenyl porphyrin chlorides with benzonitrile: formation of Rh porphyrin arene and imine complexes, 80

Electron transfer
Activation of a coordinated alkyne by electron transfer: crystal structures of [Pd{PPh$_2$CH=C(Bu^t)NN=C(Bu^t)CH$_2$PPh$_2$}-{C(CO$_2$Me)=CH(CO$_2$Me)}] and [Pd{(*Z*,*Z*)PPh$_2$CH$_2$C(Bu^t)=NN=C(Bu^t)CH$_2$PPh$_2$}{C(CO$_2$-Me)=C(CO$_2$Me)}], 318
Electrophilic aromatic substitution
Bromination and nitration reactions of metallated (Ru and Os) multiaromatic ligands and crystal structures of selected products, 262
Electrophilic substitution
Interaction of bis(trimethylsilyl)acetylene complex of titanocene with tris(pentafluorophenyl)borane. Synthesis and structure of a new type of zwitterionic metallocene (η^5-C$_5$H$_5$){η^5-[C$_5$H$_4$B(C$_6$F$_5$)$_3$]}Ti, 243
Emission spectroscopy
Crystal structures and spectroscopic properties of platina-β-diketones — a new mode of stacking in platinum complexes, 286
Enantiomeric purity
Use of *cis*-dichloro[(*S*)-α-methylbenzylamine](ethylene)Pt(II) as a chiral derivatizing agent for the determination by ^{195}Pt-NMR of the enantiomeric composition of unsaturated ethers or alcohols having a quaternary chiral carbon atom, 174
Enynyl
Alkyne insertion reactions of [RuH(κ^2-S$_2$CNEt$_2$)(CO)(PPh$_3$)$_2$]: synthesis of alkenyl, alkynyl and enynyl complexes, 20

Ferrocene
New 1,1′- or 1,2- or 1,3-bis(diphenylphosphino)ferrocenes, 365
Synthesis and properties of some novel ferrocene macrocyclic dioxopolyamines, 42
Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254
Ferrocenyl
Synthesis, structure and coordination chemistry of mono- and bis-heterocyclic-ferrocenyl derivatives, 248
Ferrocenyl ligand
Platinum(II) and palladium(II) compounds derived from [(η^5-C$_5$H$_5$)Fe{(η^5-C$_5$H$_4$)–CH=N–CH$_2$–CH$_2$–OH}], 87
Ferrocenyl ligands
Ferrocenyl ligands in mixed valence high-oxidation state rhenium complexes, 108
β-Ferrocenylvinylazide
P,*N*-Heterodifunctional ligands by selective Staüdinger reaction of β-substituted vinylazides with (*Z*)-1,2-bis(diphenylphosphanyl)ethene and formation of cyclometalled complexes of palladium(II) of these ligands — crystal and molecular structure of a new chiral cyclometallaphosphoraniminophosphane of palladium(II), 329
Fluoroalkylated carbenes
Fluoroalkylated *N*-heterocyclic carbene complexes of palladium, 409

Fluorophenyl α-amino acids
Water soluble phosphines: Part XIII. Chiral phosphine ligands with amino acid moieties, 116
Fluorophenyl boron compounds
(Fluoroorgano)fluoroboranes and -fluoroborates I: synthesis and spectroscopic characterization of potassium fluoroaryltrifluoroborates and fluoroaryldifluoroboranes, 127
Fluxional
Isolation of new polynuclear aluminium amides; trinuclear [Al-Cl(MeNCH$_2$CH$_2$NMe)$_2$(Al(NMe$_2$)Cl)(Al(NMe$_2$)$_2$)] and fluxional adamantane-like [Al$_4$Cl$_2$(NMe$_2$)$_6$(NMe)O], 36

Germaethene
Addukte der Ethene Me$_2$E=C(SiMe$_3$)$_2$ (E = Si, Ge, Sn) mit LiR und RN$_3$: Wie rasch bilden sie sich?, 304
Dimere der Ethene Me$_2$E=C(SiMe$_3$)$_2$ (E = Si, Ge, Sn): Auf welchem Wege entstehen sie aus Me$_2$EX–CM(SiMe$_3$)$_2$? Wie sind sie strukturiert?, 292
Germatranes
Preparation of germatranyl triflates: reactions of germatranes N(CH$_2$CHRO)$_3$X (X = Br, OTf, OSiMe$_3$; R = H, Me) with lithium reagents, 387
Gold
Synthesis, characterization and solution studies on heteronuclear iron–copper clusters containing diphosphine ligands: crystal structure of the dicluster [{Fe$_2$(μ-CO)(CO)$_6$(μ-PPh$_2$)-Cu}$_2$(dppp)], 215

Half-sandwich complex
Non-bridged amido cyclopentadienyl complexes of titanium: synthesis, characterization, and olefin polymerization catalysis, 179
Haloalkyls
Synthesis and crystal structures of the monomeric organotelluriu-m(IV) trihalides: *trans*-2-ethoxy-cyclohexyl-tellurium(IV) trichloride, trichloro(2-chlorobicyclo[2.2.1]hept-7-yl)-λ^4-tellurane, and mesityltellurium(IV) tribromide, 49
Haloaryls
Synthesis and crystal structures of the monomeric organotelluriu-m(IV) trihalides: *trans*-2-ethoxy-cyclohexyl-tellurium(IV) trichloride, trichloro(2-chlorobicyclo[2.2.1]hept-7-yl)-λ^4-tellurane, and mesityltellurium(IV) tribromide, 49
Heteroarylferrocene
Conversion of tributylstannylferrocene to a variety of heteroaryl ferrocenes, 348
Heteroaryltrimethyltins
Organotin mediated nitration in heteroaromatic series using tetranitromethane or dinitrogen tetroxide, 187
Heterocyclic
Synthesis, structure and coordination chemistry of mono- and bis-heterocyclic-ferrocenyl derivatives, 248
Heterocyclic bromides
Conversion of tributylstannylferrocene to a variety of heteroaryl ferrocenes, 348
Heterometallic
Synthesis, characterization and solution studies on heteronuclear iron–copper clusters containing diphosphine ligands: crystal structure of the dicluster [{Fe$_2$(μ-CO)(CO)$_6$(μ-PPh$_2$)Cu}$_2$(dppp)], 215
Heteronuclear coordination polymer
Cyano-bridged extended heteronuclear supramolecular architectures with hexacyanoferrates(II) as building blocks, 63
Heteropolyoxometalates
Monoorganotin–polyoxometal compounds: Part 1. Synthesis and crystal structure of [(^{i}PrSn)$_{11}$(V^{IV}O)O$_{14}$(OH)$_6$]Cl·2DMF·H$_2$O. The first example in a series of monoorganotin–polyoxometal compounds with icosahedral metal arrangement and various contents of tin and vanadium, 381
Hg(CF$_3$)$_2$ compound
A theoretical study of gas-phase and solid-state Hg(CF$_3$)$_2$, 374
Honeycomb structure
Cyano-bridged extended heteronuclear supramolecular architectures with hexacyanoferrates(II) as building blocks, 63
Hydride
Alkyne insertion reactions of [RuH(κ^2-S$_2$CNEt$_2$)(CO)(PPh$_3$)$_2$]: synthesis of alkenyl, alkynyl and enynyl complexes, 20
Hydrogen bond
Cyano-bridged extended heteronuclear supramolecular architectures with hexacyanoferrates(II) as building blocks, 63
Syntheses of cationic four-coordinated aluminum complexes: counter-ion induced aggregation through the N–H···I hydrogen bonds, 13

Imido
Syntheses of titanium μ-arylimido and μ-pyridylimido complexes bearing (un)substituted cyclopentadienyl ligand, 339
Indirect nuclear spin coupling constants
Implementation of the IPPP–CLOPPA–INDO/S method for the study of indirect nuclear spin coupling constants and its application to molecules containing tin nuclei, 193
Indium antimonide
Preparation of [Et$_2$InSb(SiMe$_3$)$_2$]$_3$; a trimeric single-source precursor to indium antimonide, 182
Insertionen
Addukte der Ethene Me$_2$E=C(SiMe$_3$)$_2$ (E = Si, Ge, Sn) mit LiR und RN$_3$: Wie rasch bilden sie sich?, 304
Dimere der Ethene Me$_2$E=C(SiMe$_3$)$_2$ (E = Si, Ge, Sn): Auf welchem Wege entstehen sie aus Me$_2$EX–CM(SiMe$_3$)$_2$? Wie sind sie strukturiert?, 292
Intramolecular coordination
Syntheses of titanium μ-arylimido and μ-pyridylimido complexes bearing (un)substituted cyclopentadienyl ligand, 339
Intramolecular transformation
Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254
Ionization potentials
A theoretical study of gas-phase and solid-state Hg(CF$_3$)$_2$, 374
IPPP–CLOPPA–INDO/S
Implementation of the IPPP–CLOPPA–INDO/S method for the study of indirect nuclear spin coupling constants and its application to molecules containing tin nuclei, 193
Iron
Ferrocenyl ligands in mixed valence high-oxidation state rhenium complexes, 108
Ironpentacarbonyl
Carbonyl insertion into zirconium–nitrogen bonds; synthesis and X-ray structure of a carbene complex composed of [Zr(NMe$_2$)$_4$]$_2$ and three Fe(CO)$_5$ units containing Fe→Zr donor–acceptor interactions, 403
Isocyanide
Development of a T-joint for covalent molecular construction based on 2,2′-bipyridine and phenanthroline isocyanide metal complexes, 55

Ligand
The synthesis and characterisation of bis(phenylpyridylphosphino)ethane, 103
Lithium reagents
Preparation of germatranyl triflates: reactions of germatranes N(CH$_2$CHRO)$_3$X (X = Br, OTf, OSiMe$_3$; R = H, Me) with lithium reagents, 387

Macrocyclic dioxopolyamine
Synthesis and properties of some novel ferrocene macrocyclic dioxopolyamines, 42
Manganese compounds
Geometrical and electronic structures of the acetyl complex $Re(bpy)(CO)_3(COCH_3)$ and of $[M(bpy)(CO)_4](OTf)$, M = Mn,Re, 136
Mesityl
Mesitylpseudohalogermanes $Mes_2Ge(CN)_2$ and Mes_3GeX (X = CN, NCS, N_3, NCO and OH; Mes = 2,4,6-trimethylphenyl): syntheses, crystal and molecular structures, 276
Metal amides
Carbonyl insertion into zirconium–nitrogen bonds; synthesis and X-ray structure of a carbene complex composed of $[Zr(NMe_2)_4]_2$ and three $Fe(CO)_5$ units containing $Fe \rightarrow Zr$ donor–acceptor interactions, 403
Metal complexes
Development of a T-joint for covalent molecular construction based on 2,2′-bipyridine and phenanthroline isocyanide metal complexes, 55
Metallocane conformation
Synthesis, characterization and X-ray structure of stannocanes substituted with a cyclic dithiophosphate ligand $X(CH_2CH_2S)_2Sn^nBu$ [$S_2P(OCH_2C(Et_2)CH_2O)$], (X = O, S): a study about the conformational tendencies and the relationship with the anomeric effect of the stannocane rings, 160
α-Methylated
Water soluble phosphines: Part XIII. Chiral phosphine ligands with amino acid moieties, 116
Molecular conformation
Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254
Molecular structures
Mesitylpseudohalogermanes $Mes_2Ge(CN)_2$ and Mes_3GeX (X = CN, NCS, N_3, NCO and OH; Mes = 2,4,6-trimethylphenyl): syntheses, crystal and molecular structures, 276
Molybdenum
Development of a T-joint for covalent molecular construction based on 2,2′-bipyridine and phenanthroline isocyanide metal complexes, 55
Multifunctional ligand
Syntheses, characterization and structures of chromium group carbonyl complexes containing a multifunctional $Ph_2P(o\text{-}C_6H_4)CH{=}N(CH_2)_2(o\text{-}C_6H_4N)$ ligand, 353

Nanocrystalline
Preparation of $[Et_2InSb(SiMe_3)_2]_3$: a trimeric single-source precursor to indium antimonide, 182
Niobium
Synthesis and reactivity of new oxo alkyl or oxo acyl niobocene complexes and crystal structure of $Cp'_2Nb(=O)(OC(O)CF_3)$ ($Cp' = \eta^5\text{-}C_5H_4SiMe_3$), 167
Nitration
Organotin mediated nitration in heteroaromatic series using tetranitromethane or dinitrogen tetroxide, 187
2-Nitrobenzo[*b*]furan
Organotin mediated nitration in heteroaromatic series using tetranitromethane or dinitrogen tetroxide, 187
2-Nitrobenzo[*b*]thiophene
Organotin mediated nitration in heteroaromatic series using tetranitromethane or dinitrogen tetroxide, 187
2-Nitroindoles
Organotin mediated nitration in heteroaromatic series using tetranitromethane or dinitrogen tetroxide, 187
2-Nitropyridine
Organotin mediated nitration in heteroaromatic series using tetranitromethane or dinitrogen tetroxide, 187
NMR
(Fluoroorgano)fluoroboranes and -fluoroborates I: synthesis and spectroscopic characterization of potassium fluoroaryltrifluoroborates and fluoroaryldifluoroboranes, 127
On some reactions of undecamethylcyclohexasilanyl-substituted silanes, 202
Synthesis and decomposition of bis(alkoxyethynyl)dimethyltin, 211
Use of *cis*-dichloro[(*S*)-α-methylbenzylamine](ethylene)Pt(II) as a chiral derivatizing agent for the determination by ^{195}Pt-NMR of the enantiomeric composition of unsaturated ethers or alcohols having a quaternary chiral carbon atom, 174
Nucleophilic phosphination
Water soluble phosphines: Part XIII. Chiral phosphine ligands with amino acid moieties, 116

Organogermanium
Mesitylpseudohalogermanes $Mes_2Ge(CN)_2$ and Mes_3GeX (X = CN, NCS, N_3, NCO and OH; Mes = 2,4,6-trimethylphenyl): syntheses, crystal and molecular structures, 276
ortho-Carborane–dithiolato complexes
The 16-electron dithiolene complexes (*p-cymene*)$M[S_2C_2(B_{10}H_{10})]$ (M = Ru, Os) containing both η^6-(*p-cymene*) and η^2-(*ortho*-carborane-dithiolate): adduct formation with Lewis bases, and X-ray crystal structures of (*p-cymene*)-$Ru[S_2C_2(B_{10}H_{10})](L)$ (L = PPh_3) and $\{(p\text{-}cymene)\text{-}Ru[S_2C_2(B_{10}H_{10})]\}_2(\mu\text{-}LL)$ (LL = $Ph_2PCH_2CH_2PPh_2$ and N_2H_4), 142
Orthometallation
Preparation of some dinuclear rhodium complexes with the orthometallated ligand $[2,6\text{-}(PPh_2CH_2)_2C_6H_3]^-$ and their catalytic activity for polymerization of phenylacetylene, 228
ortho-Metallation
Bromination and nitration reactions of metallated (Ru and Os) multiaromatic ligands and crystal structures of selected products, 262
Osmium
Bromination and nitration reactions of metallated (Ru and Os) multiaromatic ligands and crystal structures of selected products, 262
The 16-electron dithiolene complexes (*p-cymene*)$M[S_2C_2(B_{10}H_{10})]$ (M = Ru, Os) containing both η^6-(*p-cymene*) and η^2-(*ortho*-carborane-dithiolate): adduct formation with Lewis bases, and X-ray crystal structures of (*p-cymene*)-$Ru[S_2C_2(B_{10}H_{10})](L)$ (L = PPh_3) and $\{(p\text{-}cymene)Ru[S_2C_2(B_{10}H_{10})]\}_2(\mu\text{-}LL)$ (LL = $Ph_2PCH_2CH_2PPh_2$ and N_2H_4), 142

Palladium
Platinum(II) and palladium(II) compounds derived from $[(\eta^5\text{-}C_5H_4)Fe\{(\eta^5\text{-}C_5H_4)\text{-}CH{=}N\text{-}CH_2\text{-}CH_2\text{-}OH\}]$, 87
P,N-Heterodifunctional ligands by selective Staüdinger reaction of β-substituted vinylazides with (*Z*)-1,2-bis(diphenylphosphanyl)ethene and formation of cyclometalled complexes of palladium(II) of these ligands — crystal and molecular structure of a new chiral cyclometallaphosphoraniminophosphane of palladium(II), 329
Reactivity of cyclometallated semicarbazone complexes of Pd(II): crystal and molecular structures of $[Pd\{C_6H_4C(Et){=}NN(H)C(=O)NH_2\}(PPh_3)][ClO_4]$ and $[Pd\{C_6H_4C(Et){=}NN(H)C(=O)NH_2\}\{(Ph_2PCH_2CH_2)_2PPh\text{-}P,P,P\}][Cl]$, 1

Synthesis of the diphenylacetylene-based, tetra-amine ligand 1,2-bis(3,5-bis[(dimethylamino)methyl]phenyl)acetylene by palladium-catalysed cross-coupling: isolation and crystal structure of the catalyst *trans*-(3,5-bis[(dimethylamino)-methyl]phenyl)bis(triphenylphosphine)palladium(II) iodide, 24

Synthesis, structure and coordination chemistry of mono- and bis-heterocyclic-ferrocenyl derivatives, 248

Palladium complexes

Fluoroalkylated *N*-heterocyclic carbene complexes of palladium, 409

Palladium(II)

Influence of phenyl ring substituents in the cyclometallation of Schiff base ligands: crystal and molecular structures of [Pd-{3,4-(OCH_2O)C_6H_2C(H)=N(Cy)-*C*2,*N*}(μ-O_2CMe)]$_2$ and [Pd-{3,4-(OCH_2CH_2O)C_6H_2C(H)=N(Cy)-*C*6,*N*}(μ-O_2CMe)]$_2$, 71

p-Cymene complexes

The 16-electron dithiolene complexes (*p*-*cymene*)M[$S_2C_2(B_{10}H_{10})$] (M = Ru, Os) containing both η^6-(*p*-*cymene*) and η^2-(*ortho*-carborane-dithiolate): adduct formation with Lewis bases, and X-ray crystal structures of (*p*-*cymene*)Ru-[$S_2C_2(B_{10}H_{10})$](L) (L = PPh_3) and {(*p*-*cymene*)Ru-[$S_2C_2(B_{10}H_{10})$]}$_2$(μ-LL) (LL = $Ph_2PCH_2CH_2PPh_2$ and N_2H_4), 142

Phenanthroline

Development of a T-joint for covalent molecular construction based on 2,2′-bipyridine and phenanthroline isocyanide metal complexes, 55

Phosphine

New 1,1′- or 1,2- or 1,3-bis(diphenylphosphino)ferrocenes, 365

Preparation of some dinuclear rhodium complexes with the orthometallated ligand [2,6-(PPh_2CH_2)$_2C_6H_3$]$^-$ and their catalytic activity for polymerization of phenylacetylene, 228

Phosphine derivatives

Water soluble phosphines: Part XIII. Chiral phosphine ligands with amino acid moieties, 116

Phosphine ligands

2-Thioanisyldichlorophosphine, new starting material for the preparation of multidentate phosphine ligands: syntheses and characterization of derivatives of 2-anisyl- and 2-thioanisyldichlorophosphines, 235

Platinum

Crystal structures and spectroscopic properties of platina-β-diketones — a new mode of stacking in platinum complexes, 286

Platinum(II) and palladium(II) compounds derived from [(η^5-C_5H_5)Fe{(η^5-C_5H_4)–CH=N–CH_2–CH_2–OH}], 87

Platinum(II)

Influence of phenyl ring substituents in the cyclometallation of Schiff base ligands: crystal and molecular structures of [Pd-{3,4-(OCH_2O)C_6H_2C(H)=N(Cy)-*C*2,*N*}(μ-O_2CMe)]$_2$ and [Pd-{3,4-(OCH_2CH_2O)C_6H_2C(H)=N(Cy)-*C*6,*N*}(μ-O_2CMe)]$_2$, 71

P,*N*-Difunctional ligands

P,*N*-Heterodifunctional ligands by selective Staüdinger reaction of β-substituted vinylazides with (*Z*)-1,2-bis(diphenylphosphanyl)ethene and formation of cyclometalled complexes of palladium(II) of these ligands — crystal and molecular structure of a new chiral cyclometallaphosphoraniminophosphane of palladium(II), 329

^{31}P-NMR spectroscopy

2-Thioanisyldichlorophosphine, new starting material for the preparation of multidentate phosphine ligands: syntheses and characterization of derivatives of 2-anisyl- and 2-thioanisyldichlorophosphines, 235

Polymerization acetylene

Preparation of some dinuclear rhodium complexes with the orthometallated ligand [2,6-(PPh_2CH_2)$_2C_6H_3$]$^-$ and their catalytic activity for polymerization of phenylacetylene, 228

Polynuclear

Isolation of new polynuclear aluminium amides; trinuclear [Al-Cl(MeNCH_2CH_2NMe_2)$_2$(Al(NMe_2)Cl)(Al(NMe_2)$_2$)] and fluxional adamantane-like [Al_4Cl_2(NMe_2)$_6$(NMeO)], 36

Precursor

Preparation of [Et_2InSb($SiMe_3$)$_2$]$_3$; a trimeric single-source precursor to indium antimonide, 182

Propargyl complexes

Reactions of transition-metal η^1-propargyl complexes with sulfur dioxide and sulfur trioxide: transition-metal–carbon bond cleaving reactions of the cycloadducts which yield 1,2-oxathiole-2-oxides and -2,2-dioxides, 150

Pseudohalides

Mesitylpseudohalogermanes $Mes_2Ge(CN)_2$ and Mes_2GeX (X = CN, NCS, N_3, NCO and OH; Mes = 2,4,6-trimethylphenyl): syntheses, crystal and molecular structures, 276

Pyridine

The synthesis and characterisation of bis(phenylpyridylphosphino)ethane, 103

Pyridyl

Syntheses of titanium μ-arylimido and μ-pyridylimido complexes bearing (un)substituted cyclopentadienyl ligand, 339

Pyridylphosphines

The synthesis and characterisation of bis(phenylpyridylphosphino)ethane, 103

Rearrangement

Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254

Recyclization

Synthesis and some chemical properties of 3-ferrocenyl-3-isopropylcyclopropene: 3-ferrocenyl-3-isopropylstructures of cyclopropene and its adducts with 1,3-diphenylisobenzofuran, 254

Redox process

Activation of a coordinated alkyne by electron transfer: crystal structures of [Pd{PPh_2CH–C(Bu′)NN=C(Bu′)$CH_2$$PPh_2$}-{C($CO_2$Me)=CH($CO_2$Me)}] and [Pd{(*Z*,*Z*)$PPh_2CH_2$C(Bu′)=NN=C(Bu′)$CH_2PPh_2$}{C($CO_2$-Me)=C($CO_2$Me)}], 318

relative Reaktionsgeschwindigkeiten

Addukte der Ethene Me_2E=C($SiMe_3$)$_2$ (E = Si, Ge, Sn) mit LiR und RN_3: Wie rasch bilden sie sich?, 304

Relativistic effects

Implementation of the IPPP–CLOPPA–INDO/S method for the study of indirect nuclear spin coupling constants and its application to molecules containing tin nuclei, 193

Resolution

Water soluble phosphines: Part XIII. Chiral phosphine ligands with amino acid moieties, 116

Rhenium

Development of a T-joint for covalent molecular construction based on 2,2′-bipyridine and phenanthroline isocyanide metal complexes, 55

Ferrocenyl ligands in mixed valence high-oxidation state rhenium complexes, 108

Rhenium compounds

Geometrical and electronic structures of the acetyl complex Re(bpy)(CO)$_4$($COCH_3$) and of [M(bpy)(CO)$_4$](OTf), M = Mn,Re, 136

Rhodium
Preparation of some dinuclear rhodium complexes with the orthometallated ligand $[2,6\text{-}(PPh_2CH_2)_2C_6H_3]^-$ and their catalytic activity for polymerization of phenylacetylene, 228
Rhodium porphyrin
Diverse reactivity of rhodium β-(tetraphenyl)tetraphenyl porphyrin chlorides with benzonitrile: formation of Rh porphyrin arene and imine complexes, 80
Röntgenstrukturanalysen
Dimere der Ethene $Me_2E{=}C(SiMe_3)_2$ (E = Si, Ge, Sn): Auf welchem Wege entstehen sie aus $Me_2EX{-}CM(SiMe_3)_2$? Wie sind sie strukturiert?, 292
Ruthenium
Alkyne insertion reactions of $[RuH(\kappa^2\text{-}S_2CNEt_2)(CO)(PPh_3)_2]$: synthesis of alkenyl, alkynyl and enynyl complexes, 20
Bromination and nitration reactions of metallated (Ru and Os) multiaromatic ligands and crystal structures of selected products, 262
Reactions of metalloalkynes: Part 6. Synthesis of open tetrametallic clusters. Reaction of $[Ru(CO)_4(ethene)]$ with ethyne-1,2-diyl compounds. Single crystal X-ray structural determinations of $[Ru_4(\mu_4\text{-}CC)(\eta\text{-}C_5H_4R)_2(\mu\text{-}CO)_2(CO)_8]$ (R = H, Me), 28
The 16-electron dithiolene complexes (*p-cymene*)$M[S_2C_2(B_{10}H_{10})]$ (M = Ru, Os) containing both η^6-(*p-cymene*) and η^2-(*ortho*-carborane-dithiolate): adduct formation with Lewis bases, and X-ray crystal structures of (*p-cymene*)$Ru[S_2C_2(B_{10}H_{10})](L)$ (L = PPh_3) and $\{$(*p-cymene*)$Ru[S_2C_2(B_{10}H_{10})]\}_2(\mu\text{-}LL)$ (LL = $Ph_2PCH_2CH_2PPh_2$ and N_2H_4), 142

Schiff bases
Influence of phenyl ring substituents in the cyclometallation of Schiff base ligands: crystal and molecular structures of $[Pd\text{-}\{3,4\text{-}(OCH_2O)C_6H_2C(H){=}N(Cy)\text{-}C2,N\}(\mu\text{-}O_2CMe)]_2$ and $[Pd\text{-}\{3,4\text{-}(OCH_2CH_2O)C_6H_2C(H){=}N(Cy)\text{-}C6,N\}(\mu\text{-}O_2CMe)]_2$, 71
Semicarbazone
Reactivity of cyclometallated semicarbazone complexes of Pd(II): crystal and molecular structures of $[Pd\{C_6H_4C(Et){=}NN(H)C(=O)NH_2\}(PPh_3)][ClO_4]$ and $[Pd\{C_6H_4C(Et){=}NN(H)C(=O)NH_2\}\{(Ph_2PCH_2CH_2)_2PPh\text{-}P,P,P\}][Cl]$, 1
Silaethene
Addukte der Ethene $Me_2E{=}C(SiMe_3)_2$ (E = Si, Ge, Sn) mit LiR und RN_3: Wie rasch bilden sie sich?, 304
Dimere der Ethene $Me_2E{=}C(SiMe_3)_2$ (E = Si, Ge, Sn): Auf welchem Wege entstehen sie aus $Me_2EX{-}CM(SiMe_3)_2$? Wie sind sie strukturiert?, 292
Silene dimerization
Head-to-head versus head-to-tail dimerizations of transient silenes — the generation and dimerization behavior of 2-(2-dimethylaminoaryl)-1,1-bis(trimethylsilyl)silenes, 395
Silenes
Head-to-head versus head-to-tail dimerizations of transient silenes — the generation and dimerization behavior of 2-(2-dimethylaminoaryl)-1,1-bis(trimethylsilyl)silenes, 395
Silicon
Direct displacement of chlorine or iodine in reactions of $(Me_3Si)_3CSiRR'X$ with metal salts, 222
Head-to-head versus head-to-tail dimerizations of transient silenes — the generation and dimerization behavior of 2-(2-dimethylaminoaryl)-1,1-bis(trimethylsilyl)silenes, 395
Sonogashira coupling
Synthesis of the diphenylacetylene-based, tetra-amine ligand 1,2-bis(3,5-bis[(dimethylamino)methyl]phenyl)acetylene by palladium-catalysed cross-coupling: isolation and crystal structure of the catalyst *trans*-(3,5-bis[(dimethylamino)-methyl]phenyl)bis(triphenylphosphine)palladium(II) iodide, 24
Stannaethene
Addukte der Ethene $Me_2E{=}C(SiMe_3)_2$ (E = Si, Ge, Sn) mit LiR und RN_3: Wie rasch bilden sie sich?, 304
Dimere der Ethene $Me_2E{=}C(SiMe_3)_2$ (E = Si, Ge, Sn): Auf welchem Wege entstehen sie aus $Me_2EX{-}CM(SiMe_3)_2$? Wie sind sie strukturiert?, 292
Stannocanes
Synthesis, characterization and X-ray structure of stannocanes substituted with a cyclic dithiophosphate ligand $X(CH_2CH_2S)_2Sn^nBu$ $[S_2P(OCH_2C(Et)(CH_2O)]$, (X = O, S): a study about the conformational tendencies and the relationship with the anomeric effect of the stannocane rings, 160
Steric Hindrance
Direct displacement of chlorine or iodine in reactions of $(Me_3Si)_3CSiRR'X$ with metal salts, 222
Stille coupling
Conversion of tributylstannylferrocene to a variety of heteroaryl ferrocenes, 348
Styrene polymerization
Non-bridged amido cyclopentadienyl complexes of titanium: synthesis, characterization, and olefin polymerization catalysis, 179
Sulfur heterocycles
Reactions of transition-metal η^1-propargyl complexes with sulfur dioxide and sulfur trioxide: transition-metal–carbon bond cleaving reactions of the cycloadducts which yield 1,2-oxathiole-2-oxides and -2,2-dioxides, 150
Syntheses
Mesitylpseudohalogermanes $Mes_2Ge(CN)_2$ and Mes_3GeX (X = CN, NCS, N_3, NCO and OH; Mes = 2,4,6-trimethylphenyl): syntheses, crystal and molecular structures, 276
Synthesis
Heteroleptic diphenylamido complexes of ytterbium: syntheses and molecular structures of $(MeC_5H_4)_2YbNPh_2(THF)$ and $(C_5Me_5)_2YbNPh_2$, 359
Synthesis and properties of some novel ferrocene macrocyclic dioxopolyamines, 42

Tellurium
Synthesis and crystal structures of the monomeric organotelluriu-m(IV) trihalides: *trans*-2-ethoxy-cyclohexyl-tellurium(IV) trichloride, trichloro(2-chlorobicyclo[2.2.1]hept-7-yl)-λ^4-tellurane, and mesityltellurium(IV) tribromide, 49
Tetra-amide
Synthesis and crystal structure of $[U(\eta\text{-}C_8H_8)]_2[\mu\text{-}\eta^4\text{:}\eta^4\text{-}HN(CH_2)_3N(CH_2)_2N(CH_2)_3NH]$, a dinuclear compound with a bridging tetra-amide ligand, 313
Tetraamine ligand
Synthesis of the diphenylacetylene-based, tetra-amine ligand 1,2-bis(3,5-bis[(dimethylamino)methyl]phenyl)acetylene by palladium-catalysed cross-coupling: isolation and crystal structure of the catalyst *trans*-(3,5-bis[(dimethylamino)-methyl]phenyl)bis(triphenylphosphine)palladium(II) iodide, 24
Tetranuclear
Reactions of metalloalkynes: Part 6. Synthesis of open tetrametallic clusters. Reaction of $[Ru(CO)_4(ethene)]$ with ethyne-1,2-diyl compounds. Single crystal X-ray structural determinations of $[Ru_4(\mu_4\text{-}CC)(\eta\text{-}C_5H_4R)_2(\mu\text{-}CO)_2(CO)_8]$ (R = H, Me), 28
Tin
Monoorganotin–polyoxometal compounds: Part 1. Synthesis and crystal structure of $[(^iPrSn)_{11}(V^{IV}O)O_{14}(OH)_8]Cl{\cdot}2DMF{\cdot}H_2O$. The first example in a series of monoorganotin–polyoxometal compounds with icosahedral metal arrangement and various contents of tin and vanadium, 381

Synthesis and decomposition of bis(alkoxyethynyl)dimethyltin, 211

Titanium
Interaction of bis(trimethylsilyl)acetylene complex of titanocene with tris(pentafluorophenyl)borane. Synthesis and structure of a new type of zwitterionic metallocene (η^5-C_5H_5){η^5-[$C_5H_4B(C_6F_5)_3$]}Ti, 243
Non-bridged amido cyclopentadienyl complexes of titanium: synthesis, characterization, and olefin polymerization catalysis, 179
Syntheses of titanium μ-arylimido and μ-pyridylimido complexes bearing (un)substituted cyclopentadienyl ligand, 339

trans-Metallation
Bromination and nitration reactions of metallated (Ru and Os) multiaromatic ligands and crystal structures of selected products, 262

Tributylstannylferrocene
Conversion of tributylstannylferrocene to a variety of heteroaryl ferrocenes, 348

Trimer
Preparation of [$Et_2InSb(SiMe_3)_2]_3$; a trimeric single-source precursor to indium antimonide, 182

Tris(pentafluorophenyl)borane
Interaction of bis(trimethylsilyl)acetylene complex of titanocene with tris(pentafluorophenyl)borane. Synthesis and structure of a new type of zwitterionic metallocene (η^5-C_5H_5){η^5-[$C_5H_4B(C_6F_5)_3$]}Ti, 243

Trisyl
Direct displacement of chlorine or iodine in reactions of $(Me_3Si)_3CSiRR'X$ with metal salts, 222

Unsaturated alcohols
Use of *cis*-dichloro[(*S*)-α-methylbenzylamine](ethylene)Pt(II) as a chiral derivatizing agent for the determination by ^{195}Pt-NMR of the enantiomeric composition of unsaturated ethers or alcohols having a quaternary chiral carbon atom, 174

Unsaturated ethers
Use of *cis*-dichloro[(*S*)-α-methylbenzylamine](ethylene)Pt(II) as a chiral derivatizing agent for the determination by ^{195}Pt-NMR of the enantiomeric composition of unsaturated ethers or alcohols having a quaternary chiral carbon atom, 174

Uranium
Synthesis and crystal structure of [U(η-C_8H_8)]$_2$[μ-η^4:η^4-$HN(CH_2)_3N(CH_2)_2N(CH_2)_3NH$], a dinuclear compound with a bridging tetra-amide ligand, 313

UV–vis spectrometry
Synthesis and properties of some novel ferrocene macrocyclic dioxopolyamines, 42

Vanadium
Monoorganotin–polyoxometal compounds: Part 1. Synthesis and crystal structure of $[(^tPrSn)_{11}(V^{IV}O)O_{14}(OH)_6]Cl\cdot 2DMF\cdot H_2O$. The first example in a series of monoorganotin–polyoxometal compounds with icosahedral metal arrangement and various contents of tin and vanadium, 381

X-ray
Synthesis of the diphenylacetylene-based, tetra-amine ligand 1,2-bis(3,5-bis[(dimethylamino)methyl]phenyl)acetylene by palladium-catalysed cross-coupling: isolation and crystal structure of the catalyst *trans*-(3,5-bis[(dimethylamino)-methyl]phenyl)bis(triphenylphosphine)palladium(II) iodide, 24
The 16-electron dithiolene complexes (*p-cymene*)M[$S_2C_2(B_{10}H_{10})$] (M = Ru, Os) containing both η^6-(*p-cymene*) and η^2-(*ortho*-carborane-dithiolate): adduct formation with Lewis bases, and X-ray crystal structures of (*p-cymene*)Ru-[$S_2C_2(B_{10}H_{10})$](L) (L = PPh_3) and {(*p-cymene*)Ru-[$S_2C_2(B_{10}H_{10})$]}$_2$(μ-LL) (LL = $Ph_2PCH_2CH_2PPh_2$ and N_2H_4), 142

X-ray structure
Development of a T-joint for covalent molecular construction based on 2,2′-bipyridine and phenanthroline isocyanide metal complexes, 55
Heteroleptic diphenylamido complexes of ytterbium: syntheses and molecular structures of $(MeC_5H_4)_2YbNPh_2(THF)$ and $(C_5Me_5)_2YbNPh_2$, 359
Interaction of bis(trimethylsilyl)acetylene complex of titanocene with tris(pentafluorophenyl)borane. Synthesis and structure of a new type of zwitterionic metallocene (η^5-C_5H_5){η^5-[$C_5H_4B(C_6F_5)_3$]}Ti, 243
On some reactions of undecamethylcyclohexasilanyl-substituted silanes, 202
Synthesis and crystal structures of the monomeric organotelluriu-m(IV) trihalides: *trans*-2-ethoxy-cyclohexyl-tellurium(IV) trichloride, trichloro(2-chlorobicyclo[2.2.1]hept-7-yl)-λ^4-tellurane, and mesityltellurium(IV) tribromide, 49
Synthesis, characterization and X-ray structure of stannocanes substituted with a cyclic dithiophosphate ligand $X(CH_2zCH_2S)_2Sn^nBu$ [$S_2P(OCH_2C(Et_2)CH_2O)$], (X = O, S): a study about the conformational tendencies and the relationship with the anomeric effect of the stannocane rings, 160

X-ray structure analysis
P,N-Heterodifunctional ligands by selective Staüdinger reaction of β-substituted vinylazides with (*Z*)-1,2-bis(diphenylphosphanyl)ethene and formation of cyclometalled complexes of palladium(II) of these ligands — crystal and molecular structure of a new chiral cyclometallaphosphoraniminophosphane of palladium(II), 329

X-ray structures
Bromination and nitration reactions of metallated (Ru and Os) multiaromatic ligands and crystal structures of selected products, 262
Fluoroalkylated *N*-heterocyclic carbene complexes of palladium, 409
Water soluble phosphines: Part XIII. Chiral phosphine ligands with amino acid moieties, 116

Ytterbocene amides
Heteroleptic diphenylamido complexes of ytterbium: syntheses and molecular structures of $(MeC_5H_4)_2YbNPh_2(THF)$ and $(C_5Me_5)_2YbNPh_2$, 359

(*Z*)-1,2-bis(Diphenylphosphanyl)ethene
P,N-Heterodifunctional ligands by selective Staüdinger reaction of β-substituted vinylazides with (*Z*)-1,2-bis(diphenylphosphanyl)ethene and formation of cyclometalled complexes of palladium(II) of these ligands — crystal and molecular structure of a new chiral cyclometallaphosphoraniminophosphane of palladium(II), 329

Zerovalent complexes
Activation of a coordinated alkyne by electron transfer: crystal structures of [Pd{PPh2CH=C(Bu^t)NN=C(Bu^t)CH$_2$PPh$_2$}-{C(CO$_2$Me)=CH(CO$_2$Me)}] and [Pd{(*Z,Z*)PPh$_2$CH$_2$C(Bu^t)=NN=C(Bu^t)CH$_2$PPh$_2$}{C(CO$_2$-Me)=C(CO$_2$Me)}], 318

Zirconium
Carbonyl insertion into zirconium–nitrogen bonds; synthesis and X-ray structure of a carbene complex composed of $[Zr(NMe_2)_4]_2$ and three $Fe(CO)_5$ units containing Fe → Zr donor–acceptor interactions, 403

Zirconium–iron interaction
Carbonyl insertion into zirconium–nitrogen bonds; synthesis and X-ray structure of a carbene complex composed of $[Zr(NMe_2)_4]$, and three $Fe(CO)_5$ units containing $Fe \rightarrow Zr$ donor–acceptor interactions, 403

Zwitterionic metallocenes
Interaction of bis(trimethylsilyl)acetylene complex of titanocene with tris(pentafluorophenyl)borane. Synthesis and structure of a new type of zwitterionic metallocene $(\eta^5\text{-}C_5H_5)\{\eta^5\text{-}[C_5H_4B(C_6F_5)_3]\}Ti$, 243